arXiv:1107.3960v1 [stat.ME] 20 Jul 2011

# A new semi-parametric family of probability distributions for survival analysis

**Damien Bousquet**[*]
Laboratoire de Biostatistiques, Épidémiologie, et de Santé Publique
IURC, Université de Montpellier 1

**Jean-Pierre Daurès**[†]
Laboratoire de Biostatistiques, Épidémiologie, et de Santé Publique
IURC, Université de Montpellier 1

**Jean-Michel Marin**[‡]
Institut de Mathématiques et de Modélisation de Montpellier
Université de Montpellier 2

## Abstract

In the context of survival analysis, Marshall & Olkin (1997) introduced families of distributions by adding a scalar parameter to a given survival function, parameterized or not. In that paper, we generalize their approach. We show how it is possible to add more than a single parameter to a given distribution. We then introduce very flexible families of distributions for which we calculate some moments. Notably, we give some tractable expressions of these moments when the given baseline distribution is Log-logistic. Finally, we demonstrate how to generate sample from these new families.

KEYWORDS: survival analysis; semi-parametric probability distributions; Log-logistic and Weibull distributions.

---

[*] DAMIEN.BOUSQUET@INSERM.FR

[†] JEAN-PIERRE.DAURES@INSERM.FR

[‡] place Eugène Bataillon, CC051, 34095 Montpellier cedex 5, France, JEAN-MICHEL.MARIN@UNIV-MONTP2.FR (corresponding author)

# 1 Introduction

By various methods, new parameters can be introduced to expand families of probability distributions. This is an important issue in survival analysis (Lawless, 2003; Lee & Wang, 2003; Marshall & Olkin, 2007). For instance, although the Weibull distribution is often described as flexible, its hazard function is restricted to being monotonically increasing or monotonically decreasing, or constant. Typically, in survival analysis, the limitations of standard distributions led naturally to interest in developing extended distributions by adding further parameters to a given distribution. The approach of Marshall & Olkin (1997) works as follows. Let $S_0(x)$ denotes the survival function of a random variable, the corresponding Marshall-Olkin extended distribution has survival function,

$$\frac{aS_0(x)}{1-(1-a)S_0(x)}$$

where $a > 0$ is an added scalar parameter. These distributions have been used in many areas:

- climatology: Biondi, Kozubowski, Panorska & Saito (2008);
- hydrology: Jose, Naik & Ristić (2008);
- insurance and finance: García, Gómez-Déniz & Vázquez-Polo (2010), Jayakumar & Mathew (2008) and Kozubowski & Panorska (2008);
- medicine: Economou & Caroni (2007), Ghitany, Al-Awadhi & Alkhalfan (2007), Ghitany, Al-Hussaini & Al-Jarallah (2005), Gómez-Déniz (2009), Gupta & Peng (2009) and Jose, Naik & Ristić (2008);
- engineering: Adamidis, Dimitrakopoulou & Loukas (2005), Adamidis & Loukas (1998), Gupta, Lvin, & Peng (2010), Prabhakar Murthy, Bulmer & Eccleston (2004), Silva, Barreto-Souza & Cordeiro (2010) and Zhang & Xie (2007)...

For a review on the use of the extended Marshall and Olkin distributions, one can see Nadarajah (2008). There is also a lot of methodological works around the Marshall and Olkin extended distributions. Independently of Marshall & Olkin (1997), Adamidis & Loukas (1998) introduced the same type of extended distributions but only in the baseline Weibull case. In Adamidis, Dimitrakopoulou & Loukas (2005), it is shown that the Marshall and Olkin extended distributions can be viewed as continuous mixtures. The baseline exponential case has been investigated by Silva, Barreto-Souza & Cordeiro (2010) and Srinivasa-Rao, Ghitany & Kantam (2009). For the more general baseline Weibull case, one can see Ghitany, Al-Hussaini & Al-Jarallah (2005); Zhang & Xie (2007); Gupta, Lvin, & Peng (2010).

In Sankaran & Jayakumar (2008), it is shown that the Marshall and Olkin extended distributions satisfy the property of proportional odds functions. This property has been used notably by Economou & Caroni (2007), Caroni (2008) and Gupta & Peng (2009) to introduce covariates.

Some authors have investigated the behavior of the Marshall and Olkin extended distributions when the baseline distributions are Lomax, Ghitany, Al-Awadhi & Alkhalfan (2007), Burr, and Pareto, Jayakumar & Mathew (2008), $q$-Weibull, Jose, Naik & Ristić (2008) and normal García, Gómez-Déniz & Vázquez-Polo (2010) distributions. Moreover, using the Marshall and Olkin approach, Gómez-Déniz (2009) introduced a generalization of the discrete Geometric distribution.

In the baseline exponential case, Kozubowski & Panorska (2008) highlighted the link between the truncated logistic distribution and the Marshall and Olkin extended one.

Finally, some authors have investigated the generalization to the multivariate case, one can see for instance: Thomas & Jose (2004), Sankaran & Jayakumar (2008), Jose, Ristić & Joseph (2009), Yeh (2009) and Yeh (2010). Random minima and maxima related to Marshall-Olkin distributions have received attention in the litterature, Arnold (1996), Biondi, Kozubowski, Panorska & Saito (2008), Ghitany, Al-Awadhi & Alkhalfan (2007), Jose, Ristić & Joseph (2009), Thomas & Jose (2004), Yeh (2009) and Yeh (2010).



Figure 1: Hazard rate curve of a standard Weibull distribution with parameter $(2, 2)$ (left), of an extended Marshall and Olkin distribution with added parameter $a = 0.05$ (baseline Weibull $(2, 2)$) (center) and of an extended distribution obtained using our approach: two parameters ($a_1 = 10^{-6}$ and $a_2 = 0.15$) are added to a baseline Weibull $(2, 2)$ (right).

In that paper, we generalize the approach of Marshall and Olkin in order to obtain more flexible distributions. Indeed, coming back to baseline Weibull one, it is easy to show that the hazard rate curve of the extended Marshall and Olkin distribution has at most two waves. As shown in Figure 1, using our approach, we can obtain more rich hazard rate curve. Our proposal is introduced in Section 2. In Section 3, we investigate some properties of these extended distributions for various baseline cases.

## 2 The new semi-parametric family

In Marshall & Olkin (1997), it is introduced a new semi-parametric family of probability distributions. One real parameter $a > 0$ is added to a given univariate probability distribution $F_0 : \mathbb{R} \to [0, 1]$ by using an increasing function denoted by $g_a$. Indeed, the Marshal and Olkin extended probability distribution is given by

$$g_a\left(F_0(x)\right) = \frac{F_0(x)}{a + (1 - a)F_0(x)} \,.$$

Here, we generalize this support function with an arbitrary number of external parameters.

Let $S_0 = 1 - F_0$ and $q \in \mathbb{N}^*$. Clearly, for any $(a_1, \ldots, a_q) \in \,]0, +\infty[^q$ and any $u \in [0, 1]$, we have,

$$\left\{ \sum_{i=1}^{q} a_i - \left( \sum_{i=1}^{q} a_i - q \right) u \right\} \neq 0 \,.$$

We define the function $g_{a_1,\ldots,a_q} : [0, 1] \to \mathbb{R}$ by

$$g_{a_1,\ldots,a_q}(u) = q^q \frac{u \displaystyle\prod_{i=2}^{q} (a_i + u - a_i u)}{\left\{ \displaystyle\sum_{i=1}^{q} a_i - \left( \sum_{i=1}^{q} a_i - q \right) u \right\}^q} \,.$$

As a generalization of the Marshal and Olkin proposal, we define the Let us now define the functions $F_{a_1,\ldots,a_q}$ and $S_{a_1,\ldots,a_q}$ such that :

$$\begin{array}{rccl} F_{a_1,\ldots,a_q} : & \mathbb{R} & \to & \mathbb{R} \\ & x & \mapsto & g_{a_1,\ldots,a_q}\{F_0(x)\} = q^q \frac{F_0(x) \prod_{i=2}^{q} \{a_i + F_0(x) - a_i F_0(x)\}}{\left\{\sum_{i=1}^{q} a_i - (\sum_{i=1}^{q} a_i - q) F_0(x)\right\}^q} \; ; \end{array}$$

**Theorem 2.1** *For any integer $q \in \mathbb{N}^*$ and any $(a_1, \ldots, a_q) \in \,]0, +\infty[^q$, the application $F_{a_1,\ldots,a_q}$ from $\mathbb{R}$ to $\mathbb{R}$ takes its values in $[0, 1]$, and the function $F_{a_1,\ldots,a_q}$ from $\mathbb{R}$ to $[0, 1]$ defines a probability distribution.*

**Proof**

The proof is given in Appendix. □

If we replace in Theorem 2.1, the cumulative probability function $F_0$ by the survival function $S_0$, we still obtain a survival function. This defines a new probability distribution. In that work, we only consider the initial construction.

For any $a > 0$, the function $F_{a,\dots,a}$ from $\mathbb{R}$ to $[0,1]$ is a Marshall-Olkin extended probability distribution. That is, for any real number $x$, we have,

$$F_{a,\dots,a}(x) = \frac{F_0(x)}{a + (1-a)F_0(x)} \,,$$

or, in another way,

$$S_{a,\dots,a}(x) = \frac{aS_0(x)}{1 - (1-a)S_0(x)} \,.$$

Let us now introduce the following lemma.

**Lemma 2.1** *We suppose that there exists* $t \in \mathbb{R}$ *such that* $0 < F_0(t) < 1$.
*In that case, there exists a unique interval* $J$ *of* $\mathbb{R}$ *such that, for any* $x \in (\mathbb{R} - J) \cap ]-\infty, t]$ *we have* $F_0(x) = 0$*, for any* $x \in (\mathbb{R} - J) \cap [t, +\infty[$ *we have* $F_0(x) = 1$*, and for any* $x$ *in* $J$ *we have* $0 < F_0(x) < 1$.

The proof of this lemma is trivial. The interval $J$ does not correspond to the usual definition of the support of a probability distribution (the smallest closed set whose complement has probability zero). We use that specific definition for technical reasons.

Let us suppose that $F_0$ is continuous on $\mathbb{R}$ and $C^1$ on $J$, the interval $J$ is defined by Lemma 2.1. Let $f_0$ be the probability density function of $F_0$. We can take $f_0$ as,

$$\begin{array}{rccl} f_0 : & \mathbb{R} & \rightarrow & \mathbb{R}_+ \\ & x & \mapsto & F_0'(x) \quad \text{if } x \in J \\ & & & 0 \quad \text{otherwise} \end{array}$$

Here, it is very easy to verify that for any $q \in \mathbb{N}^*$ and any $(a_1, \dots, a_q) \in ]0, +\infty[^q$, the function $F_{a_1,\dots,a_q}$ from $\mathbb{R}$ into $[0,1]$ is continuous on $\mathbb{R}$ and $C^1$ on $J$.

Moreover, the probability density function $f_{a_1,\ldots,a_q}$ of $F_{a_1,\ldots,a_q}$ can be taken as,

$$\begin{array}{rccll} f_{a_1,\ldots,a_q}: & \mathbb{R} & \to & \mathbb{R}_+ & \\ & x & \mapsto & F'_{a_1,\ldots,a_q}(x) = g'_{a_1,\ldots,a_q}\left\{F_0(x)\right\} f_0(x) & \text{if } x \in J \\ & & & 0 & \text{otherwise} \end{array}$$

For instance if $q = 2$, we get for any $x \in J$,

$$f_{a_1,a_2}(x) = 4\frac{a_1 a_2 F_0(x) + a_1^2 F_0(x) - a_1 a_2 - a_1^2 - 2a_2 F_0(x)}{\left\{a_1 F_0(x) + a_2 F_0(x) - a_1 - a_2 - 2F_0(x)\right\}^3} f_0(x)\,.$$

Let us now introduce two Lemmas.

**Lemma 2.2** *Let $q \in \mathbb{N}^*$ and $a_1, \ldots, a_q$ be strictly positive real numbers such that $a_1 + \cdots + a_q > \frac{q}{2}$. For any $u \in [-1, 1]$,*

$$\sum_{i=1}^{q} a_i - \left(\sum_{i=1}^{q} a_i - q\right) u \neq 0\,,$$

*and*

$$g_{a_1,\ldots,a_q}(u) = \sum_{m=1}^{\infty} c_{q,a_1,\ldots,a_q,m} u^m$$

*where, for any $m \in \mathbb{N}^*$,*

$$c_{q,a_1,\ldots,a_q,m} =$$

$$\frac{q^q}{(q-1)!}\frac{\prod_{i=2}^{q} a_i}{\left(\sum_{i=1}^{q} a_i\right)^q}\left\{\sum_{j+k=m,1\leq j\leq q} \sigma_{j-1}\,(k+q-1)\cdots(k+1)\left(\frac{\sum_{i=1}^{q} a_i - q}{\sum_{i=1}^{q} a_i}\right)^k\right\}$$

*with*

$$\sigma_i = \left\{\begin{array}{ll} 1 & \text{if} \quad i = 0 \\ \sum_{2\leq j_1<\ldots<j_i\leq q} \frac{1-a_{j_1}}{a_{j_1}} \times \cdots \times \frac{1-a_{j_i}}{a_{j_i}} & \text{if} \quad 1 \leq i \leq q-1 \\ 0 & \text{if} \quad i \geq q \end{array}\right\}.$$

*Note that we have*

$$\sum_{m=1}^{\infty} c_{q,a_1,\ldots,a_q,m} = 1\,.$$

**Proof**

The proof is given in Appendix. □

We trivially deduce the following corollary used in the proof of Theorem 2.2.

**Corollary 2.1** *Let $q \in \mathbb{N}^*$ and $a_1, \ldots, a_q$ be strictly positive real numbers such that $a_1 + \cdots + a_q \geq q$ and $a_i \leq 1$ for $2 \leq i \leq q$. Then, for any $m \in \mathbb{N}^*$,*

$$c_{q,a_1,\ldots,a_q,m} \geq 0\,.$$

**Lemma 2.3** *Let $q \in \mathbb{N}^*$ and $a_1, \ldots, a_q$ be strictly positive real numbers such that $a_1 + \cdots + a_q < 2q$. For any $u \in [0,2]$,*

$$\sum_{i=1}^{q} a_i - \left(\sum_{i=1}^{q} a_i - q\right) u \neq 0\,,$$

*and*

$$g_{a_1,\ldots,a_q}(u) = 1 + \sum_{m=1}^{\infty} d_{q,a_1,\ldots,a_q,m}(u-1)^m$$

*where, for $m \in \mathbb{N}^*$,*

$$\begin{aligned} d_{q,a_1,\ldots,a_q,m} = \quad & \tfrac{1}{(q-1)!}\Big\{(m+q-1)\cdots(m+1)\big(\tfrac{\sum_{i=1}^{q} a_i - q}{q}\big)^m \\ & + \textstyle\sum_{k+j=m,1\leq j\leq q}(k+q-1)\cdots(k+1)\big(\tfrac{\sum_{i=1}^{q} a_i - q}{q}\big)^k(\sigma_j + \sigma_{j-1})\Big\} \end{aligned}$$

*with*

$$\sigma_i = \left\{ \begin{array}{ll} 1 & \text{if} \quad i = 0 \\ \sum_{2 \leq j_1 < \ldots < j_i \leq q} (1 - a_{j_1}) \times \cdots \times (1 - a_{j_i}) & \text{if} \quad 1 \leq i \leq q-1 \\ 0 & \text{if} \quad i \geq q \end{array} \right\}.$$

**Proof**

The proof is given in Appendix. □

We now study the expectation of our family of distributions when the baseline probability distribution $F_0$ is continuous on $\mathbb{R}$ and $C^1$ on $J$.

**Theorem 2.2** *Let* $(\Omega, \mathcal{A}, P)$ *be a probability space,* $q \in \mathbb{N}^*$, $(a_1, \ldots, a_q) \in ]0, +\infty[^q$, $X_0 : \Omega \to \mathbb{R}$ *and* $X_{a_1,\ldots,a_q} : \Omega \to \mathbb{R}$ *be random variables such that* $X_0 \sim F_0$ *and* $X_{a_1,\ldots,a_q} \sim F_{a_1,\ldots,a_q}$ *and* $w : \mathbb{R} \to \mathbb{R}$ *be a borelian function.*
*If* $F_0$ *is continuous on* $\mathbb{R}$ *and* $C^1$ *on* $J$ *and if* $E(|w \circ X_0|) = \int_{\mathbb{R}} |w(x)| f_0(x) dx \in \mathbb{R}_+$, *then*

$$E(|w \circ X_{a_1,\ldots,a_q}|) \in \mathbb{R}_+ .$$

*Moreover, if* $a_1, \ldots, a_q$ *be strictly positive real numbers such that* $a_1 + \cdots + a_q \geq q$ *and* $a_i \leq 1$ *for any* $2 \leq i \leq q$, *then*

$$E(|w \circ X_{a_1,\ldots,a_q}|) \leq a_1 E(|w \circ X_0|) .$$

**Proof**

The proof is given in Appendix. □

As a direct consequence, if $a_1, \dots, a_q$ be strictly positive real numbers such that $a_1 + \cdots + a_q \geq q$ and $a_i \leq 1$ for any $2 \leq i \leq q$ and, $\int_{\mathbb{R}} |x|^r f_0(x)dx \in \mathbb{R}_+$ for a real number $r$, then

$$E(|X_{a_1,\dots,a_q}|^r) \leq a_1 E(|X_0|^r)\,.$$

The following proposition will be used to derive explicit formulae on some expectations when the baseline probability distribution is a classical Log-logistic.

**Proposition 2.1** *We suppose that $F_0$ is bijective from $J$ to $]0,1[$. Let $(\Omega, \mathcal{A}, P)$ be a probability space, $(a_1, a_2) \in \, ]0, +\infty[^2$ such that $a_1 \neq a_2$, $X_{a_1,a_2} : \Omega \to \mathbb{R}$ and $V_{a_1,a_2} : \Omega \to \mathbb{R}$ be independent random variables such that $X_{a_1,a_2} \sim F_{a_1,a_2}$ and $V_{a_1,a_2} \sim \mathrm{Exp}\left(\frac{a_1+a_2}{|a_1-a_2|}\right)$.*
*Define for any $\omega \in \Omega$ such that $X_{a_1,a_2}(\omega) \in J$,*

$$Y_{a_1,a_2}(\omega) = \log\left[\frac{1}{F_0\{X_{a_1,a_2}(\omega)\}} - 1\right]\,,$$

*and for any $\omega \in \Omega$ such that $X_{a_1,a_2}(\omega) \in \mathbb{R} - J$,*

$$Y_{a_1,a_2}(\omega) = 0\,.$$

*Then the random variable,*

$$L_{a_1,a_2} = \begin{cases} Y_{a_1,a_2} + V_{a_1,a_2} - \log\frac{2}{a_1+a_2} & \text{if} \quad a_1 > a_2 \\ -Y_{a_1,a_2} + V_{a_1,a_2} - \log\frac{a_1+a_2}{2} & \text{if} \quad a_1 < a_2 \end{cases}$$

*is distributed according to a logistic distribution.*

**Proof**

Let us denote by $\alpha_{a_1,a_2}$ the cumulative probability function of $V_{a_1,a_2}$, by $r_{a_1,a_2}$ the one of $Y_{a_1,a_2}$, by $t_{a_1,a_2}$ the one of $-Y_{a_1,a_2}$, and by $u_{a_1,a_2}$ the one of $L_{a_1,a_2}$.

For any $u \in \mathbb{R}$, we have

$$r_{a_1,a_2}(u) = 1 - g_{a_1,a_2}\left\{\frac{1}{1+\exp(u)}\right\}$$

and

$$t_{a_1,a_2}(u) = g_{a_1,a_2}\left\{\frac{\exp(u)}{1+\exp(u)}\right\} .$$

We first suppose that $a_1 > a_2$ and for any $(u,t) \in \mathbb{R}^2$, we define

$$\theta_{a_1,a_2,u}(t) = -\frac{2(a_1+a_2)\exp(u+\frac{2a_2}{a_2-a_1}t)}{\{2\exp(t)+(a_1+a_2)\exp(u)\}^2} .$$

For any $(u,t) \in \mathbb{R} \times [0,+\infty[$, we clearly have

$$\theta'_{a_1,a_2,u}(t) = r'_{a_1,a_2}(u-t)\alpha'_{a_1,a_2}(t)$$

and

$$\int_{\mathbb{R}_+} r'_{a_1,a_2}(u-t)\alpha'_{a_1,a_2}(t)dt = \frac{2(a_1+a_2)\exp(u)}{\{2+(a_1+a_2)\exp(u)\}^2} .$$

Therefore, for any $v \in \mathbb{R}$,

$$\int_{-\infty}^{v}\int_{\mathbb{R}_+} r'_{a_1,a_2}(u-t)\alpha'_{a_1,a_2}(t)dtdu = \frac{(a_1+a_2)\exp(v)}{2+(a_1+a_2)\exp(v)} = u_{a_1,a_2}\left(v - \log\frac{2}{a_1+a_2}\right)$$

and

$$u_{a_1,a_2}(v) = \frac{\exp(v)}{1+\exp(v)} .$$

We now suppose that $a_2 > a_1$ and for any $(u,t) \in \mathbb{R}^2$ and we define

$$\theta_{a_1,a_2,u}(t) = -\frac{2(a_1+a_2)\exp(u - \frac{2a_1}{a_2-a_1}t)}{\{2\exp(u) + (a_1+a_2)\exp(t)\}^2} .$$

For any $(u,t) \in \mathbb{R} \times [0,+\infty[$, we clearly have

$$\theta'_{a_1,a_2,u}(t) = t'_{a_1,a_2}(u-t)\alpha'_{a_1,a_2}(t)$$

and

$$\int_{\mathbb{R}_+} t'_{a_1,a_2}(u-t)\alpha'_{a_1,a_2}(t)dt = \frac{2(a_1+a_2)\exp(u)}{\{2\exp(u) + a_1 + a_2\}^2} .$$

Therefore, for any $v \in \mathbb{R}$,

$$\int_{-\infty}^{v} \int_{\mathbb{R}_+} t'_{a_1,a_2}(u-t)\alpha'_{a_1,a_2}(t)dtdu = \frac{2\exp(v)}{2\exp(v) + a_1 + a_2} = u_{a_1,a_2}\left(v - \log\frac{a_1+a_2}{2}\right)$$

and

$$u_{a_1,a_2}(v) = \frac{\exp(v)}{1+\exp(v)} .$$

$\square$

In Marshall & Olkin (1997), it is shown a specific property of the introduced semi-parametric family: the proposed distributions are geometric extreme stable. That is, they are both minimum and maximum stable, when the random indexation variable of the sample size is distributed according to a geometric distribution. We now consider maximum stability for our semi-parametric family.

**Proposition 2.2** *Let $q \in \mathbb{N}^*$ and $a_1, \ldots, a_q$ be strictly positive real numbers such that $a_1 + \cdots + a_q \geq q$ and $a_i \leq 1$ for any $2 \leq i \leq q$. Let $(\Omega, \mathcal{A}, P)$ be a probability space and $N_{a_1,\ldots,a_q} : \Omega \to \mathbb{N}^*$ be a random variable such that, for any $m \in \mathbb{N}^*$,*

$$\mathrm{pr}(N_{a_1,\ldots,a_q} = m) = c_{q,a_1,\ldots,a_q,m} \,.$$

- *If, for $i \in \mathbb{N}^*$, $X_i : \Omega \to \mathbb{R}$ be mutually independent random variables, independent from $N_{a_1,\ldots,a_q}$, such that $X_i \sim F_0$, then*

$$V_{a_1,\ldots,a_q} = \max(X_1, \ldots, X_{N_{a_1,\ldots,a_q}}) \sim F_{a_1,\ldots,a_q} \,.$$

- *If, for $b > 0$ and $i \in \mathbb{N}^*$, $X_i : \Omega \to \mathbb{R}$ be mutually independent random variables, independent from $N_{a_1,\ldots,a_q}$ such that $X_i \sim F_{b,\ldots,b}$, then*

$$V_{a_1,\ldots,a_q} = \max(X_1, \ldots, X_{N_{a_1,\ldots,a_q}}) \sim F_{ba_1,\ldots,ba_q} \,.$$

**Proof**

Part one: we have for any real number $x$,

$$\mathrm{pr}(V_{a_1,\ldots,a_q} \leq x) = \sum_{m=1}^{\infty} \mathrm{pr}(N_{a_1,\ldots,a_q} = m) F_0(x)^m \,,$$

$$\mathrm{pr}(V_{a_1,\ldots,a_q} \leq x) = g_{a_1,\ldots,a_q}\left\{F_0(x)\right\} \,,$$

$$\mathrm{pr}(V_{a_1,\ldots,a_q} \leq x) = F_{a_1,\ldots,a_q}(x) \,.$$

Part two: for any $(u, a_1, \ldots, a_q, b) \in [0,1] \times ]0, +\infty[^{q+1}$, we have,

$$g_{a_1,\ldots,a_q}\left\{g_{b,\ldots,b}(u)\right\} = g_{ba_1,\ldots,ba_q}(u)\,.$$

□

More generally for $q \geq 2$, the new probability distributions are not maximum stable.

# 3 Some properties of the extended distributions

## 3.1 $F_0$ is Weibull

We first suppose that $F_0(x) = (1 - \exp(-x))\,\mathbb{I}_{x\geq 0}$ (exponential distribution with expectation equal to 1).

**Lemma 3.1** *Let* $F_0(x) = (1 - \exp(-x))\,\mathbb{I}_{x\geq 0}$ *and* $f_0$ *is the corresponding probability density function. For any* $r > 0$ *and any* $m \in \mathbb{N}^*$*, we have*

$$\int_{\mathbb{R}} x^r F_0(x)^{m-1} f_0(x)dx = r\Gamma(r)\sum_{j=0}^{m-1}\binom{m-1}{j}(-1)^j(j+1)^{-r-1}\,.$$

**Proof**

Let $r > 0$, $p \in \mathbb{N}$ and $z \geq 0$,

$$F_0(z)^p = \sum_{j=0}^{p} \binom{p}{j} (-1)^j \exp(-jz)$$

and

$$z^r F_0(z)^p f_0(z) = \sum_{j=0}^{p} \binom{p}{j} (-1)^j z^r \exp\left\{-(j+1)z\right\} .$$

Then,

$$\int_{\mathbb{R}_+} x^r F_0(x)^p f_0(x) dx = \sum_{j=0}^{p} \binom{p}{j} (-1)^j \int_{\mathbb{R}_+} x^r \exp\left\{-(j+1)x\right\} dx$$

and

$$\int_{\mathbb{R}_+} x^r F_0(x)^p f_0(x) dx = \sum_{j=0}^{p} \binom{p}{j} (-1)^j \frac{r\Gamma(r)}{(j+1)^{r+1}} = r\Gamma(r) \sum_{j=0}^{p} \binom{p}{j} (-1)^j (j+1)^{-r-1} .$$

$\square$

**Proposition 3.1** *Let* $F_0(x) = (1 - \exp(-x))\, \mathbb{I}_{x \geq 0}$ *and* $X_{a_1,\ldots,a_q} \sim F_{a_1,\ldots,a_q}$.

*If* $a_1, \ldots, a_q$ *be strictly positive real numbers such that* $a_1 + \cdots + a_q \geq q$ *and* $a_i \leq 1$ *for* $2 \leq i \leq q$, $E({X_{a_1,\ldots,a_q}}^r) = r\Gamma(r) \sum_{m=1}^{\infty} m c_{q,a_1,\ldots,a_q,m} \sum_{j=0}^{m-1} \binom{m-1}{j} (-1)^j (j+1)^{-r-1}$ *for any* $r > 0$.

*If* $a_1, \ldots, a_q$ *be strictly positive real numbers such that* $q \leq a_1 + \cdots + a_q < 2q$ *and* $a_i \leq 1$ *for* $2 \leq i \leq q$, $E({X_{a_1,\ldots,a_q}}^r) = r\Gamma(r) \sum_{m=1}^{\infty} (-1)^{m-1} d_{q,a_1,\ldots,a_q,m} m^{-r}$ *for any* $r > 0$.

**Proof**

The proof of this result is almost trivial using Lemmas 2.2, 2.3 and 3.1. $\square$

We now consider the case of the classical Weibull distribution with two parameters, that is

$$F_0(x) = F_0^{b_1,b_2}(x) = \left(1 - \exp\left\{-\left(\frac{x}{b_1}\right)^{b_2}\right\}\right) \mathbb{I}_{x \geq 0}$$

with $(b_1, b_2) \in ]0, +\infty[^2$. Let $X_{a_1,\ldots,a_q}^{b_1,b_2} \sim F_{a_1,\ldots,a_q}^{b_1,b_2}$ ($F_{a_1,\ldots,a_q}^{b_1,b_2}$ be the corresponding parameter augmented distribution). We can easily verify that $X_{a_1,\ldots,a_q}^{b_1,b_2} = b_1 {X_{a_1,\ldots,a_q}^{1,1}}^{\frac{1}{b_2}}$ in distribution and, then for any $r > 0$,

$$E\left({X_{a_1,\ldots,a_q}^{b_1,b_2}}^{r}\right) = {b_1}^{r} E\left({X_{a_1,\ldots,a_q}^{1,1}}^{\frac{r}{b_2}}\right) .$$

Finally, we consider the case of the generalized Weibull distribution with three parameters, that is

$$F_0^{b_1,b_2,b_3}(x) = \left(1 - \exp\left[1 - \left\{1 + (\frac{x}{b_1})^{b_2}\right\}^{\frac{1}{b_3}}\right]\right) \mathbb{I}_{x \geq 0}$$

with $(b_1, b_2, b_3) \in ]0, +\infty[^3$. Let $X_{a_1,\ldots,a_q}^{b_1,b_2,b_3} \sim F_{a_1,\ldots,a_q}^{b_1,b_2,b_3}$. We can easily verify that $X_{a_1,\ldots,a_q}^{b_1,b_2,b_3} = b_1 \left\{(1 + X_{a_1,\ldots,a_q}^{1,1,1})^{b_3} - 1\right\}^{\frac{1}{b_2}}$ in distribution and then, if we suppose in addition that $m_2 = \frac{1}{b_2} \in \mathbb{N} - \{0\}$ and $b_3 \in \mathbb{N}^*$, it follows that for any integer $m \in \mathbb{N}^*$,

$$E({X_{a_1,\ldots,a_q}^{b_1,b_2,b_3}}^{m}) = b_1^m \left\{\sum_{k=0}^{mm_2} \sum_{j=0}^{b_3 k} \binom{mm_2}{k} \binom{b_3 k}{j} (-1)^{mm_2 - k} E({X_{a_1,\ldots,a_q}^{1,1,1}}^{k b_3 j})\right\} .$$

## 3.2 $F_0$ is Log-logistic

We now consider the case of the standard Log-logistic distribution, that is

$$F_0(x) = \left(\frac{x}{1+x}\right)\mathbb{I}_{x\geq 0}\,.$$

**Proposition 3.2** *Let* $F_0(x) = \left(\frac{x}{1+x}\right)\mathbb{I}_{x\geq 0}$ *and* $X_{a_1,\ldots,a_q} \sim F_{a_1,\ldots,a_q}$.

*If* $a_1,\ldots,a_q$ *be strictly positive real numbers such that* $a_1+\cdots+a_q \geq q$ *and* $a_i \leq 1$ *for* $2 \leq i \leq q$, *then for any* $r$ *such that* $|r| < 1$,

$$E({X_{a_1,\ldots,a_q}}^r) = \sum_{m=1}^{\infty} m c_{q,a_1,\ldots,a_q,m}\mathrm{Beta}(1-r, m+r)\,.$$

*If* $a_1,\ldots,a_q$ *be strictly positive real numbers such that* $q \leq a_1+\cdots+a_q < 2q$ *and* $a_i \leq 1$ *for* $2 \leq i \leq q$, *then for any* $r$ *such that* $|r| < 1$,

$$E({X_{a_1,\ldots,a_q}}^r) = \sum_{m=1}^{\infty} (-1)^{m-1} m d_{q,a_1,\ldots,a_q,m}\mathrm{Beta}(m-r, 1+r)\,.$$

**Proof**

The proof of this result is almost trivial using Lemmas 2.2 and 2.3. □

We now consider the case of the classical Log-logistic distribution with two parameters, that is

$$F_0^{b_1,b_2}(x) = \left(\frac{x^{b_2}}{{b_1}^{b_2} + x^{b_2}}\right)\mathbb{I}_{x\geq 0}$$

with $(b_1, b_2) \in ]0,+\infty[^2$.

Let $X^{b_1,b_2}_{a_1,\dots,a_q} \sim F^{b_1,b_2}_{a_1,\dots,a_q}$. As in the Weibull case, we can easily to verify that $X^{b_1,b_2}_{a_1,\dots,a_q} = b_1 {X^{1,1}_{a_1,\dots,a_q}}^{\frac{1}{b_2}}$ in distribution, and then for any $r$ such that $|r| < b_2$,

$$E({X^{b_1,b_2}_{a_1,\dots,a_q}}^r) = {b_1}^r E({X^{1,1}_{a_1,\dots,a_q}}^{\frac{r}{b_2}}) \, .$$

**Proposition 3.3** *Let* $F_0\left(\frac{x}{1+x}\right)\mathbb{I}_{x\geq 0}$ *and* $X_{a_1,a_2} \sim F_{a_1,a_2}$. *For any* $r$ *such that* $|r| < 1$,

$$E({X_{a_1,a_2}}^r) = \left(\frac{a_1+a_2}{2}\right)^r \frac{r\pi}{\sin(r\pi)} \left(\frac{a_1-a_2}{a_1+a_2} r + 1\right) \, .$$

**Proof**

When the baseline distribution is a classical Log-logistic with two parameters, the extended Marshall-Olkin distribution with one external parameter gives again a classical Log-logistic distribution. Therefore, the case $a_1 = a_2$ is trivial.

Let us now consider the case $a_1 > a_2$

Using Proposition 2.1, we have

$$\frac{a_1+a_2}{2} \exp(-L_{a_1,a_2}) = \exp(-Y_{a_1,a_2}) \exp(-V_{a_1,a_2}) \, .$$

Then

$$\left(\frac{a_1+a_2}{2}\right)^r \{\exp(-L_{a_1,a_2})\}^r = \{\exp(-Y_{a_1,a_2})\}^r \{\exp(-V_{a_1,a_2})\}^r$$

and

$$\left(\frac{a_1+a_2}{2}\right)^r E(\{\exp(-L_{a_1,a_2})\}^r) = E(\{\exp(-Y_{a_1,a_2})\}^r) E(\{\exp(-V_{a_1,a_2})\}^r) \, .$$

The random variable $\exp(-L_{a_1,a_2})$ has a Log-logistic distribution and for any $r$ such that $|r| < 1$,

$$E(\{\exp(-L_{a_1,a_2})\}^r) = \frac{r\pi}{\sin(r\pi)} \,.$$

Moreover,

$$E(\{\exp(-V_{a_1,a_2})\}^r) = \int_0^1 \frac{a_1+a_2}{a_1-a_2} x^{r+\frac{a_1+a_2}{a_1-a_2}-1} dx = \frac{a_1+a_2}{(a_1-a_2)r+(a_1+a_2)} \,.$$

Therefore,

$$E(\{\exp(-Y_{a_1,a_2})\}^r) = E({X_{a_1,a_2}}^r) = \left(\frac{a_1+a_2}{2}\right)^r \frac{r\pi}{\sin(r\pi)} \frac{(a_1-a_2)r+(a_1+a_2)}{a_1+a_2} \,.$$

We use the same type of reasoning for the case $a_1 < a_2$. $\square$

If $X_{a_1,a_2}^{b_1,b_2} \sim F_{a_1,a_2}^{b_1,b_2}$ then for any $r$ such that $|r| < b_2$

$$E({X_{a_1,a_2}^{b_1,b_2}}^r) = {b_1}^r \left(\frac{a_1+a_2}{2}\right)^{\frac{r}{b_2}} \frac{r\pi}{b_2 \sin(\frac{r\pi}{b_2})} \left\{ \frac{r(a_1-a_2)}{b_2(a_1+a_2)} + 1 \right\} \,.$$

### 3.3 Random sample generation

When the distribution $F_0$ admits a density with respect to Lebesgue measure on $\mathbb{R}$, if we are able to generate a sample from $F_0$, we can use the accept-reject algorithm to generate a sample from $F_{a_1,\ldots,a_q}$.

Indeed, following the proof of Theorem 2.2, for any $(a_1, \dots, a_q) \in ]0, +\infty[^q$,

$$\begin{aligned} f_{a_1,\dots,a_q}(x) \leq \; & \left\{ \frac{\prod_{i=2}^{q} \max(1,a_i)}{\min(1,\sum_{i=1}^{q} a_i/q)^q} + |\sum_{i=1}^{q} a_i - q| \frac{\prod_{i=2}^{q} \max(1,a_i)}{\min(1,\sum_{i=1}^{q} a_i/q)^{q+1}} \right. \\ & \left. + \frac{\sum_{i=2}^{q} |1-a_i| \prod_{2 \leq j \leq q, j \neq i} \max(1,a_j)}{\min(1,\sum_{i=1}^{q} a_i/q)^q} \right\} f_0(x)\,. \end{aligned}$$

Moreover, if $a_1 + \ldots + a_q \geq q$ and if for any integer $i$ such that $2 \leq i \leq q$, we have $a_i \leq 1$, then

$$f_{a_1,\dots,a_q}(x) \leq a_1 f_0(x)\,.$$

# 4 Conclusion

We shown how to generalize the approach of Marshall & Olkin (1997) in order to obtain more flexible families. We investigated some properties of the introduced distributions. We are now working on the parameter estimation task. The preliminary results are extremely encouraging, notably the application of our proposal on some real survival datasets.

# Appendix

## Proof of Theorem 2.1

The case $q = 1$ is trivial, it corresponds to the Marshall and Olkin derivation.

### Part one:

Let us first prove that for any $(a_1, a_2) \in \,]0, +\infty[^2$, the function $g_{a_1,a_2}$ from $[0, 1]$ into $\mathbb{R}$ is increasing on $[0, 1]$ and takes values in $[0, 1]$.

For any $u \in [0, 1]$, we have,

$$g'_{a_1,a_2}(u) = \frac{4(a_1a_2u + a_1^2u - a_1a_2 - a_1^2 - 2a_2u)}{(a_1u + a_2u - a_1 - a_2 - 2u)^3} \,.$$

Moreover,

$$a_1a_2(u-1) + a_1^2(u-1) - 2a_2u < 0\,,$$

$$a_1(u-1) + a_2(u-1) - 2u < 0\,,$$

and,

$$\{a_1(u-1) + a_2(u-1) - 2u\}^3 < 0\,.$$

Therefore,

$$g'_{a_1,a_2}(u) > 0\,.$$

Thus, the function $g_{a_1,a_2}$ from $[0, 1]$ to $\mathbb{R}$ is increasing on $[0, 1]$. As $g_{a_1,a_2}(0) = 0$ and $g_{a_1,a_2}(1) = 1$, the function $g_{a_1,a_2}$ from $[0, 1]$ into $\mathbb{R}$ takes values in $[0, 1]$.

**Part two:**

Let us now prove by induction that for any integer $q$ such that $q \geq 2$ and any $(a_1, \dots, a_q) \in ]0, +\infty[^q$, the function $g_{a_1,\dots,a_q}$ from $[0,1]$ to $\mathbb{R}$ is increasing on $[0,1]$ and takes values in $[0,1]$.

For any integer $q$ such that $q \geq 1$, we denote by $(H_q)$ the condition such that for any $(a_1, \dots, a_q) \in ]0, +\infty[^q$, the function $g_{a_1,\dots,a_q}$ from $[0,1]$ to $\mathbb{R}$ is increasing on $[0,1]$ and takes values in $[0,1]$.

Let $q$ be an integer such that $q \geq 3$ and suppose that the condition $(H_{q-1})$ is verified.
In that case, for any $(u, a_1, \dots, a_q) \in [0,1] \times ]0, +\infty[^q$, we have,

$$g_{a_1,\dots,a_q}(u) = g_{\frac{a_1}{a_q},\dots,\frac{a_{q-1}}{a_q},1}\left\{g_{a_q,\dots,a_q}(u)\right\}$$

and,

$$g_{\frac{a_1}{a_q},\dots,\frac{a_{q-1}}{a_q},1}(u) = \frac{q^q u \prod_{i=2}^{q-1}(\frac{a_i}{a_q} + u - \frac{a_i}{a_q}u)}{\left[\sum_{i=1}^{q-1} \frac{a_i}{a_q} + 1 - \left\{\sum_{i=1}^{q-1} \frac{a_i}{a_q} - (q-1)\right\} u\right]^q} \,,$$

$$g_{\frac{a_1}{a_q},\dots,\frac{a_{q-1}}{a_q},1}(u) = \frac{q^q}{(q-1)^{q-1}} g_{\frac{a_1}{a_q},\dots,\frac{a_{q-1}}{a_q}}(u) \frac{\left[\sum_{i=1}^{q-1} \frac{a_i}{a_q} - \left\{\sum_{i=1}^{q-1} \frac{a_i}{a_q} - (q-1)\right\} u\right]^{q-1}}{\left[\sum_{i=1}^{q-1} \frac{a_i}{a_q} + 1 - \left\{\sum_{i=1}^{q-1} \frac{a_i}{a_q} - (q-1)\right\} u\right]^q} \,.$$

Also,

$$\left[\sum_{i=1}^{q-1} \frac{a_i}{a_q} + 1 - \left\{\sum_{i=1}^{q-1} \frac{a_i}{a_q} - (q-1)\right\} u\right]^q = \sum_{j=0}^{q} \binom{q}{j} \left[\sum_{i=1}^{q-1} \frac{a_i}{a_q} - \left\{\sum_{i=1}^{q-1} \frac{a_i}{a_q} - (q-1)\right\} u\right]^{q-j} \,,$$

$$\frac{\left[\sum_{i=1}^{q-1} \frac{a_i}{a_q} + 1 - \left\{\sum_{i=1}^{q-1} \frac{a_i}{a_q} - (q-1)\right\} u\right]^q}{\left[\sum_{i=1}^{q-1} \frac{a_i}{a_q} - \left\{\sum_{i=1}^{q-1} \frac{a_i}{a_q} - (q-1)\right\} u\right]^{q-1}} = \sum_{j=0}^{q} \binom{q}{j} \left[\sum_{i=1}^{q-1} \frac{a_i}{a_q} - \left\{\sum_{i=1}^{q-1} \frac{a_i}{a_q} - (q-1)\right\} u\right]^{1-j} \,.$$

For any $u \in [0,1]$, let us define,

$$\eta_{a_1,\ldots,a_q}(u) = \frac{\left[\sum_{i=1}^{q-1} \frac{a_i}{a_q} + 1 - \left\{\sum_{i=1}^{q-1} \frac{a_i}{a_q} - (q-1)\right\} u\right]^q}{\left[\sum_{i=1}^{q-1} \frac{a_i}{a_q} - \left\{\sum_{i=1}^{q-1} \frac{a_i}{a_q} - (q-1)\right\} u\right]^{q-1}} .$$

We have,

$$\eta'_{a_1,\ldots,a_q}(u)$$

$$= -\left\{\sum_{i=1}^{q-1} \frac{a_i}{a_q} - (q-1)\right\} - \sum_{j=2}^{q} \binom{q}{j}(1-j)\left\{\sum_{i=1}^{q-1} \frac{a_i}{a_q} - (q-1)\right\}\left[\sum_{i=1}^{q-1} \frac{a_i}{a_q} - \left\{\sum_{i=1}^{q-1} \frac{a_i}{a_q} - (q-1)\right\} u\right]^{-j}$$

$$= -\left\{\sum_{i=1}^{q-1} \frac{a_i}{a_q} - (q-1)\right\}\left\{1 + \sum_{j=2}^{q} \binom{q}{j}(1-j)\left[\sum_{i=1}^{q-1} \frac{a_i}{a_q} - \left\{\sum_{i=1}^{q-1} \frac{a_i}{a_q} - (q-1)\right\} u\right]^{-j}\right\} ,$$

and then,

$$\eta'_{a_1,\ldots,a_q}(1) = -\left\{\sum_{i=1}^{q-1} \frac{a_i}{a_q} - (q-1)\right\}\left\{1 + \sum_{j=2}^{q} \binom{q}{j}(1-j)(q-1)^{-j}\right\} .$$

For any $x \in \mathbb{R} - \{0\}$, let us define,

$$r_q(x) = \sum_{j=2}^{q} \binom{q}{j} x^{-j+1}$$

We have

$$r_q(x) = x\left\{\left(\frac{1}{x} + 1\right)^q - 1 - \frac{q}{x}\right\} ,$$

$$r_q(x) = x\left(\frac{1}{x} + 1\right)^q - x - q \, ,$$

and,

$$r_q'(x)=\sum_{j=2}^{q}\binom{q}{j}(1-j)\,x^{-j}=\left(\frac{1}{x}+1\right)^{q}-\frac{q}{x}\left(\frac{1}{x}+1\right)^{q-1}-1=\left(\frac{1}{x}+1\right)^{q-1}\left(\frac{1-q}{x}+1\right)-1\,,$$

$$r_q'(q-1)+1=0\,.$$

Therefore,

$$\eta'_{a_1,\ldots,a_q}(1)=0\,.$$

Moreover, we have,

$$\eta''_{a_1,\ldots,a_q}(u)=\left\{\sum_{i=1}^{q-1}\frac{a_i}{a_q}-(q-1)\right\}^2\sum_{j=2}^{q}\binom{q}{j}(j-1)j\left[\sum_{i=1}^{q-1}\frac{a_i}{a_q}-\left\{\sum_{i=1}^{q-1}\frac{a_i}{a_q}-(q-1)\right\}u\right]^{-j-1}\,.$$

Then clearly, $\eta''_{a_1,\ldots,a_q}(u)\ \geq\ 0$ and $\eta'_{a_1,\ldots,a_q}(u)\ \leq\ \eta'_{a_1,\ldots,a_q}(1)\ =\ 0$. Therefore, the function $u\mapsto\frac{1}{\eta_{a_1,\ldots,a_q}(u)}$ from $[0,1]$ to $\mathbb{R}_+$ is increasing on $[0,1]$.

Using condition $(H_{q-1})$, the function $u\mapsto g_{\frac{a_1}{a_q},\ldots,\frac{a_{q-1}}{a_q}}(u)$ from $[0,1]$ to $\mathbb{R}$ is increasing and positive. It follows that the function $g_{a_1,\ldots,a_q}$ from $[0,1]$ to $\mathbb{R}_+$ is increasing on $[0,1]$.

Moreover, $g_{a_1,\ldots,a_q}(0)=0$ and $g_{a_1,\ldots,a_q}(1)=1$. Thus, the condition $(H_q)$ is verified.

Therefore, we get the result by induction.

**Part three:**

Let us now prove that the function $F_{a_1,\ldots,a_q}$ from $\mathbb{R}$ to $[0,1]$ defines a probability distribution.

The function $F_{a_1,\ldots,a_q}$ from $\mathbb{R}$ to $\mathbb{R}$ takes its values in $[0,1]$, and it is a right continuous increasing function. Moreover, $\lim_{x\to-\infty}F_{a_1,\ldots,a_q}(x)=0$ and $\lim_{x\to+\infty}F_{a_1,\ldots,a_q}(x)=1$.

Thus, it is a probability distribution.

## Proof of Lemma 2.2

Let $(a_1, \ldots, a_q) \in ]0, +\infty[^q$ such that $a_1 + \cdots + a_q > \frac{q}{2}$. For any $u \in [-1, 1]$, we have,

$$g_{a_1,\ldots,a_q}(u) = q^q \frac{u \prod_{i=2}^q \{a_i + (1-a_i)u\}}{\{\sum_{i=1}^q a_i - (\sum_{i=1}^q a_i - q)u\}^q} \, ,$$

$$\frac{1}{\{\sum_{i=1}^q a_i - (\sum_{i=1}^q a_i - q)u\}^q} = \frac{1}{(\sum_{i=1}^q a_i)^q} \times \frac{1}{\left\{1 - \frac{(\sum_{i=1}^q a_i - q)u}{\sum_{i=1}^q a_i}\right\}^q} \, ,$$

$$-1 < \frac{(\sum_{i=1}^q a_i - q)u}{\sum_{i=1}^q a_i} < 1 \, .$$

Then,

$$\frac{1}{\left\{1 - \frac{(\sum_{i=1}^q a_i - q)u}{\sum_{i=1}^q a_i}\right\}^q} = \frac{1}{(q-1)!} \sum_{k=0}^{\infty} (k+q-1) \cdots (k+1) \left\{ \frac{(\sum_{i=1}^q a_i - q)\, u}{\sum_{i=1}^q a_i} \right\}^k .$$

Moreover,

$$u \prod_{i=2}^q \{a_i + (1-a_i)u\} = (\prod_{i=2}^q a_i) u \prod_{i=2}^q (1 + \frac{1-a_i}{a_i} u) = (\prod_{i=2}^q a_i) u \prod_{i=1}^{q-1} (1 + \frac{1-a_{i+1}}{a_{i+1}} u) \, ,$$

$$u \prod_{i=2}^q \{a_i + (1-a_i)u\} = (\prod_{i=2}^q a_i) u \sum_{i=0}^{q-1} \sigma_i u^i = (\prod_{i=2}^q a_i) \sum_{i=0}^{q-1} \sigma_i u^{i+1} = (\prod_{i=2}^q a_i) \sum_{i=1}^{q} \sigma_{i-1} u^i = (\prod_{i=2}^q a_i) \sum_{j=1}^{\infty} \sigma_{j-1} u^j \, .$$

Therefore,

$$g_{a_1,\ldots,a_q}(u) = \frac{q^q}{(q-1)!} \frac{\prod_{i=2}^q a_i}{(\sum_{i=1}^q a_i)^q} (\sum_{j=1}^{\infty} \sigma_{j-1} u^j) \sum_{k=0}^{\infty} (k+q-1) \cdots (k+1) \left\{ \frac{(\sum_{i=1}^q a_i - q)\, u}{\sum_{i=1}^q a_i} \right\}^k ,$$

$$g_{a_1,\ldots,a_q}(u) = \frac{q^q}{(q-1)!} \frac{\prod_{i=2}^q a_i}{(\sum_{i=1}^q a_i)^q} \sum_{m=1}^{\infty} \left\{ \sum_{j+k=m, 1 \le j \le q} \sigma_{j-1} (k+q-1) \cdots (k+1) \left( \frac{\sum_{i=1}^q a_i - q}{\sum_{i=1}^q a_i} \right)^k \right\} u^m \, .$$

## Proof of Lemma 2.3

Let $(a_1, \dots, a_q) \in ]0, +\infty[^q$ such that $a_1 + \cdots + a_q < 2q$ and $u \in [0, 2]$. We define $w = u - 1$, we get that $w \in [-1, 1]$ and,

$$g_{a_1,\dots,a_q}(u) = q^q \frac{u \prod_{i=2}^q (a_i + u - a_i u)}{\{\sum_{i=1}^q a_i - (\sum_{i=1}^q a_i - q)u\}^q} \, ,$$

$$g_{a_1,\dots,a_q}(u) = q^q \frac{(1+w) \prod_{i=2}^q \{a_i + 1 + w - a_i(1+w)\}}{\{\sum_{i=1}^q a_i - (\sum_{i=1}^q a_i - q)(1+w)\}^q} \, ,$$

$$g_{a_1,\dots,a_q}(u) = q^q \frac{(1+w) \prod_{i=2}^q \{1 + (1 - a_i)w\}}{\{q - (\sum_{i=1}^q a_i - q)w\}^q} \, ,$$

$$g_{a_1,\dots,a_q}(u) = \frac{(1+w) \prod_{i=2}^q \{1 + (1 - a_i)w\}}{(1 - \frac{\sum_{i=1}^q a_i - q}{q} w)^q} \, .$$

Moreover, $-1 < \frac{\sum_{i=1}^q a_i - q}{q} < 1$, then $-1 < \frac{\sum_{i=1}^q a_i - q}{q} w < 1$ and,

$$\frac{1}{(1 - \frac{\sum_{i=1}^q a_i - q}{q} w)^q} = \frac{1}{(q-1)!} \sum_{k=0}^{\infty} (k + q - 1) \cdots (k+1) \left\{ \frac{(\sum_{i=1}^q a_i - q)w}{q} \right\}^k \, .$$

Also,

$$\prod_{i=2}^q \{1 + (1 - a_i)w\} = \sum_{m=0}^{q-1} \sigma_m w^m = \sum_{m=0}^{\infty} \sigma_m w^m \, ,$$

$$(1+w) \prod_{i=2}^q \{1 + (1 - a_i)w\} = \sum_{m=0}^{\infty} \sigma_m w^m + \sum_{m=0}^{\infty} \sigma_m w^{m+1} \, ,$$

$$(1+w) \prod_{i=2}^q \{1 + (1 - a_i)w\} = 1 + \sum_{m=1}^{\infty} (\sigma_m + \sigma_{m-1}) w^m \, .$$

Then,

$$g_{a_1,\dots,a_q}(u) = \frac{1}{(q-1)!} \left\{ 1 + \sum_{m=1}^{\infty} (\sigma_m + \sigma_{m-1}) w^m \right\} \left[ \sum_{k=0}^{\infty} (k + q - 1) \cdots (k+1) \left\{ \frac{(\sum_{i=1}^q a_i - q)w}{q} \right\}^k \right] \, ,$$

$$
\begin{aligned}
g_{a_1,\dots,a_q}(u) = \quad & \tfrac{1}{(q-1)!}\left\{\textstyle\sum_{k=0}^{\infty}(k+q-1)\cdots(k+1)\left(\frac{\sum_{i=1}^q a_i-q}{q}\right)^k w^k \right.\\
&\left. +\textstyle\sum_{m=1}^{\infty}\sum_{k+j=m,1\le j\le q}(k+q-1)\dots(k+1)\left(\frac{\sum_{i=1}^q a_i-q}{q}\right)^k(\sigma_j+\sigma_{j-1})w^m\right\}\,,
\end{aligned}
$$

$$
\begin{aligned}
g_{a_1,\dots,a_q}(u) = \quad & 1+\tfrac{1}{(q-1)!}\left[\textstyle\sum_{m=1}^{\infty}\left\{(m+q-1)\dots(m+1)\left(\frac{\sum_{i=1}^q a_i-q}{q}\right)^m\right.\right.\\
&\left.\left.+\textstyle\sum_{k+j=m,1\le j\le q}(k+q-1)\dots(k+1)\left(\frac{\sum_{i=1}^q a_i-q}{q}\right)^k(\sigma_j+\sigma_{j-1})\right\}w^m\right]\,.
\end{aligned}
$$

## Proof of Theorem 2.2

### Part one:

Let us prove first that for any integer $q$ such that $q \geq 1$, any $(a_1,\dots,a_q) \in \,]0,+\infty[^q$ and any $u \in [0,1]$, we have,

$$g'_{a_1,\dots,a_q}(1) = a_1\,,$$

$$g'_{a_1,\dots,a_q}(u) \geq 0\,,$$

$$g'_{a_1,\dots,a_q}(u)$$

$$\leq \frac{\prod_{i=2}^q \max(1,a_i)}{\min(1,\frac{\sum_{i=1}^q a_i}{q})^q}+|\sum_{i=1}^q a_i-q|\frac{\prod_{i=2}^q \max(1,a_i)}{\min(1,\frac{\sum_{i=1}^q a_i}{q})^{q+1}}+\frac{\sum_{i=2}^q |1-a_i|\prod_{2\le j\le q, j\neq i}\max(1,a_j)}{\min(1,\frac{\sum_{i=1}^q a_i}{q})^q}\,.$$

For any $u \in [0,1]$, we have,

$$
\begin{aligned}
g'_{a_1,\dots,a_q}(u) = \quad & q^q\tfrac{\prod_{i=2}^q(a_i+u-a_iu)}{\left\{\sum_{i=1}^q a_i-(\sum_{i=1}^q a_i-q)u\right\}^q}\\
&+q^{q+1}(\textstyle\sum_{i=1}^q a_i-q)\tfrac{u\prod_{i=2}^q(a_i+u-a_iu)}{\left\{\sum_{i=1}^q a_i-(\sum_{i=1}^q a_i-q)u\right\}^{q+1}}\\
&+q^q\tfrac{u\sum_{i=2}^q(1-a_i)\prod_{2\le j\le q,j\neq i}(a_j+u-a_ju)}{\left\{\sum_{i=1}^q a_i-(\sum_{i=1}^q a_i-q)u\right\}^q}\,.
\end{aligned}
$$

Then,

$$g'_{a_1,\dots,a_q}(1) = 1 + \sum_{i=1}^{q} a_i - q + \sum_{i=2}^{q}(1-a_i) = a_1\,.$$

For any $u \in [0,1]$, we have

$$\begin{aligned} g'_{a_1,\dots,a_q}(u) \le \quad & |q^q \tfrac{\prod_{i=2}^{q}(a_i+u-a_iu)}{\left\{\sum_{i=1}^{q} a_i-(\sum_{i=1}^{q} a_i-q)u\right\}^q}| \\ & +|q^{q+1}(\textstyle\sum_{i=1}^{q} a_i - q)\tfrac{u\prod_{i=2}^{q}(a_i+u-a_iu)}{\left\{\sum_{i=1}^{q} a_i-(\sum_{i=1}^{q} a_i-q)u\right\}^{q+1}}| \\ & +|q^q \tfrac{u\sum_{i=2}^{q}(1-a_i)\prod_{2\le j\le q, j\ne i}(a_j+u-a_ju)}{\left\{\sum_{i=1}^{q} a_i-(\sum_{i=1}^{q} a_i-q)u\right\}^q}|\,. \end{aligned}$$

Moreover, for any integer $2 \le i \le q$

$$0 \le \min(1, a_i) \le a_i + u - a_i u \le \max(1, a_i)\,,$$

$$0 < \min(q, \sum_{i=1}^{q} a_i) \le \sum_{i=1}^{q} a_i - (\sum_{i=1}^{q} a_i - q)u \le \max(q, \sum_{i=1}^{q} a_i)\,.$$

Therefore,

$$\begin{aligned} g'_{a_1,\dots,a_q}(u) \le \quad & q^q \tfrac{\prod_{i=2}^{q}\max(1,a_i)}{\min(q,\sum_{i=1}^{q} a_i)^q} \\ & +q^{q+1}|\textstyle\sum_{i=1}^{q} a_i - q|\tfrac{\prod_{i=2}^{q}\max(1,a_i)}{\min(q,\sum_{i=1}^{q} a_i)^{q+1}} \\ & +q^q \tfrac{\sum_{i=2}^{q}|1-a_i|\prod_{2\le j\le q, j\ne i}\max(1,a_j)}{\min(q,\sum_{i=1}^{q} a_i)^q}\,, \end{aligned}$$

$$g'_{a_1,\dots,a_q}(u) \le \frac{\prod_{i=2}^{q}\max(1,a_i)}{\min(1,\frac{\sum_{i=1}^{q} a_i}{q})^q} + |\sum_{i=1}^{q} a_i - q|\frac{\prod_{i=2}^{q}\max(1,a_i)}{\min(1,\frac{\sum_{i=1}^{q} a_i}{q})^{q+1}} + \frac{\sum_{i=2}^{q}|1-a_i|\prod_{2\le j\le q, j\ne i}\max(1,a_j)}{\min(1,\frac{\sum_{i=1}^{q} a_i}{q})^q}\,.$$

As the function $g_{a_1,\dots,a_q}$ from $[0,1]$ to $\mathbb{R}$ is increasing and differentiable on $[0,1]$, for any $u \in [0,1]$,

$$g'_{a_1,\dots,a_q}(u) \ge 0\,.$$

**Part two:**

Let us suppose that $\int_{\mathbb{R}} |w(x)| f_0(x)dx \in \mathbb{R}_+$. We have

$$E(|w \circ X_{a_1,\ldots,a_q}|) = \int_{\mathbb{R}} |w(x)| g'_{a_1,\ldots,a_q}\left\{F_0(x)\right\} f_0(x)dx\,.$$

Using the result given in Part one, for any $u \in [0,1]$,

$$\begin{aligned} E(|w \circ X_{a_1,\ldots,a_q}|) \leq & \left\{ \frac{\prod_{i=2}^{q} \max(1,a_i)}{\min(1,\frac{\sum_{i=1}^{q} a_i}{q})^{q}} \right. \\ & + |\textstyle\sum_{i=1}^{q} a_i - q| \frac{\prod_{i=2}^{n} \max(1,a_i)}{\min(1,\frac{\sum_{i=1}^{q} a_i}{q})^{q+1}} \\ & \left. + \frac{\sum_{i=2}^{q} |1-a_i| \prod_{2\leq j\leq q, j\neq i} \max(1,a_j)}{\min(1,\frac{\sum_{i=1}^{q} a_i}{q})^{q}} \right\} E(|w \circ X_0|)\,. \end{aligned}$$

Therefore,

$$E(|w \circ X_{a_1,\ldots,a_q}|) \in \mathbb{R}_+\,.$$

By the corollary 2.1, if $q \geq 1$ and $a_1, \ldots, a_q$ be strictly positive real numbers such that $a_1 + \cdots + a_q \geq q$ and $a_i \leq 1$ for any $2 \leq i \leq q$, then we have that for any integer $m$ which is greater than or equal to one and any $u \in [0,1]$,

$$g''_{a_1,\ldots,a_q}(u) \geq 0$$

Therefore,

$$g'_{a_1,\ldots,a_q}(u) \leq g'_{a_1,\ldots,a_q}(1)\,,$$

$$E(|w \circ X_{a_1,\ldots,a_q}|) \leq a_1 E(|w \circ X_0|)\,.$$

# References

ADAMIDIS, K., DIMITRAKOPOULOU, T. & LOUKAS, S. (2005). On an extension of the exponential-geometric distribution *Statistics & Probability Letters* **73, 3**, 259–269

ADAMIDIS, K. & LOUKAS, S. (1998). A lifetime distribution with decreasing failure rate *Statistics & Probability Letters* **39, 1**, 35–42

ARNOLD, B. C. (1996). Distributions with Logistic Marginals and/or Conditionals *Institute of Mathematical Statistics* **28**, Lecture Notes - Monograph Series

BAGDONAVICIUS, V. & NIKULIN, M. (2002). Accelerated Life Models, Modeling and Statistical Analysis *Monographs on Statistics and Applied Probability 94, Chapman & Hall/Crc*

BIONDI, F., KOZUBOWSKI, T. J., PANORSKA, A. K. & SAITO, L. (2008). A new stochastic model of episode peak and duration for eco-hydro-climatic applications *Ecological Modelling* **211, 3-4**, 383–395

CARONI, C. (2008). Testing for the Marshall-Olkin extended form of the Weibull distribution *Statistical papers*

ECONOMOU, P. & CARONI, C. (2007). Parametric proportional odds frailty models *Communications in Statistics: Simulation & Computation* **36, 4-6**, 1295–1307

GARCÍA, V. J., GÓMEZ-DÉNIZ, E., VÁZQUEZ-POLO, F. J. (2010). A new skew generalization of the normal distribution: Properties and applications *Computational Statistics & Data Analysis* **54, 8**, 2021–2034

GHITANY, M. E., AL-AWADHI, F. A. & ALKHALFAN, L. A. (2007). Marshall-Olkin extended Lomax distribution and its application to censored data *Communications in Statistics: Theory & Methods* **36, 9-12**, 1855–1866

GHITANY, M. E., AL-HUSSAINI, E. K. & AL-JARALLAH, R. A. (2005). Marshall-Olkin extended Weibull distribution and its application to censored data *Journal of Applied Statistics* **32, 10**, 1025–1034

GÓMEZ-DÉNIZ, E. (2009). Another generalization of the geometric distribution *Test*

GUPTA, R. C. & PENG, C. (2009). Estimating reliability in proportional odds ratio models *Computational Statistics & Data Analysis* **53, 4**, 1495–1510

GUPTA, R. C., LVIN, S., PENG, C. (2010). Estimating turning points of the failure rate of the extended Weibull distribution *Computational Statistics & Data Analysis* **54, 4**, 924–934

JAYAKUMAR, K. & MATHEW, T. (2008). On a generalization to Marshall-Olkin scheme and its application to Burr type XII distribution *Statistical Papers* **49**, 421–439

JOSE, K. K., NAIK, S. R. & RISTIĆ, M. M. (2008). Marshall-Olkin q-Weibull distribution and max-min processes *Statistical Papers*

JOSE, K. K., RISTIĆ, M. M. & JOSEPH, A. (2009). Marshall-Olkin bivariate Weibull distributions and processes *Statistical Papers*

KOZUBOWSKI, T. J. & PANORSKA, A. K. (2008). A Mixed Bivariate Distribution Connected with Geometric Maxima of Exponential Variables *Communications in Statistics: Theory & Methods* **37, 18**, 2903–2923

LAWLESS, J. F. (2003). Statistical models and methods for lifetime data *Wiley Series in Probability and Statistics*

LEE, E. T. & WANG, J. W. (2003). Statistical methods for survival data analysis *Wiley Series in Probability and Statistics*

MARSHALL, A. W. & OLKIN, I. (1997). A new method for adding a parameter to a family of distributions with application to the exponential and Weibull families *Biometrika* **84, 3**, 641–652

MARSHALL, A. W. & OLKIN, I. (2007). Life distributions, Structure of Nonparametric, Semiparametric, and Parametric Families *Springer Series in Statistics*

NADARAJAH, S. (2008). Marshall and Olkin's Distributions *Acta Applicandae Mathematicae* **103, 1**, 87–100

PRABHAKAR MURTHY, D. N., BULMER M., ECCLESTON, J. A. (2004). Weibull model selection for reliability modelling *Reliability Engineering & System Safety* **86, 3**, 257–267

SANKARAN, P. G. & JAYAKUMAR, K. (2008). On proportional odds models *Statistical Papers* **49**, 779–789

SILVA, R. B., BARRETO-SOUZA, W. & CORDEIRO, G. M. (2010). A new distribution with decreasing, increasing and upside-down bathtub failure rate *Computational Statistics & Data Analysis* **54, 4**, 935–944

SRINIVASA RAO, G., GHITANY, M. E. & KANTAM, R. R. L. (2009). Reliability test plans for Marshall-Olkin extended exponential distribution *Applied Mathematical Sciences* **3, 55**, 2745–2755

THOMAS, A. & JOSE, K. K. (2004). Bivariate semi-Pareto minification processes *Metrika* **59, 3**, 305–313

YEH, H-C. (2009). Multivariate semi-Weibull distributions *Journal of Multivariate Analysis* **100, 8**, 1634–1644

YEH, H-C. (2010). Multivariate semi-logistic distributions *Journal of Multivariate Analysis* **101, 4**, 893–908

ZHANG, T. & XIE, M. (2007). Failure data analysis with extended Weibull distribution *Communications in Statistics: Simulation & Computation* **36, 1-3**, 579–592